

A New Measurement of the Energy Dependence of Nuclear Transparency for Large Momentum Transfer $^{12}\text{C}(p, 2p)$ Scattering

A. Leksanov^a, J. Alster^b, G. Asryan^c, Y. Averichev^h,
 D. Barton^d, V. Baturin^{a,e}, N. Bukhtojarova^{d,e}, A. Carroll^c,
 A. Schetkovsky^{a,e}, S. Heppelmann^a, T. Kawabata^f, A. Malki^b,
 Y. Makdisi^c, E. Minina^a, I. Navon^b, H. Nicholson^g,
 A. Ogawa^a, Y. Panebratsev^h, E. Piasetzky^b, S. Shimanskiy^h,
 A. Tangⁱ, J.W. Watsonⁱ, H. Yoshida^f, D. Zhalov^a

^a*Physics Department, Pennsylvania State University, University Park, PA 16801, USA*

^b*School of Physics and Astronomy, Sackler Faculty of Exact Sciences, Tel Aviv University, Ramat Aviv 69978, Israel*

^c*Yerevan Physics Institute, Yerevan 375036, Armenia*

^d*Collider-Accelerator Department, Brookhaven National Laboratory, Upton, NY 11973, USA*

^e*Petersburg Nuclear Physics Institute, Gatchina, St. Petersburg 188350, Russia*

^f*Department of Physics, Kyoto University, Sakyo, Kyoto, 606-8502, Japan*

^g*Department of Physics, Mount Holyoke College, South Hadley, MA 01075, USA*

^h*J.I.N.R., Dubna, 141980, Russia*

ⁱ*Department of Physics, Kent State University, Kent, OH 44242, USA*

Abstract. We present a new measurement of the energy dependence of nuclear transparency from AGS experiment E850, performed using the EVA solenoidal spectrometer, upgraded since 1995. Using a secondary beam from the AGS accelerator, we simultaneously measured pp elastic scattering from hydrogen and $(p, 2p)$ quasi-elastic scattering in carbon at incoming momenta of 5.9, 8.0, 9.0, 11.7 and 14.4 GeV/c. This incident momentum range corresponds to a Q^2 region between 4.8 and 12.7 (GeV/c)². The detector allowed us to do a complete kinematic analysis for the center-of-mass polar angles in the range $85^\circ - 90^\circ$. We report on the measured variation of the nuclear transparency with energy and compare the new results with previous measurements.

Color Transparency(CT) is the predicted reduction in the initial and final state interactions, which may take place when a large transverse momentum(p_t), quasi-exclusive scattering, involving hadrons, occurs in nuclear matter. We study nuclear transparency for $(p, 2p)$ quasi-elastic scattering. It can be informally defined as

$$T_r = \frac{\frac{d\sigma}{dt}(pp \text{ quasi-elastic in nucleus})}{Z \frac{d\sigma}{dt}(pp \text{ elastic in hydrogen})} . \quad (1)$$

Theoretical predictions for the variations of this quantity with energy are very model dependent. For example, Glauber-based calculations show no energy dependence above 5 GeV with a value around thirty percent, whereas QCD based models predict an increase of T_r with energy, reaching eventually an asymptotic value of unity [1]. Interpretations of fluctuating energy dependence of the transparency [2] at the values of Q^2 above 10 (GeV/c)² have been presented in the models described in Refs. [3] and [4].

Many widely accepted pictures of CT dynamics are based on the assumption that selection of small-size Fock state configurations occurs in certain hard scattering processes and that these small configurations have a reduced cross-section for reinteraction with the surrounding nucleus. The protons remain in these configurations during a time period which is different for different energies and large on the scale of nuclear processes [5].

We performed our measurement using the EVA apparatus located in the C1 line of the AGS at BNL. Two differential Cerenkov counters were used for the incident particle identification. The beam flux measurement was performed using a scintillating-fiber beam hodoscope located in the upstream part of the apparatus. We used solid carbon and polyethylene targets, whose dimensions were 5.1 cm $\times$ 5.1 cm $\times$ 6.6 cm. The AGS provided a secondary beam with approximately $1\text{-}2 \times 10^7$ particles per 2 second spill and about 20 spills per minute. EVA is a magnetic spectrometer with tracking capabilities. It was built around a solenoidal superconducting magnet which produced nearly uniform field in the beam direction allowing for charged particle transverse momentum(p_t) measurements. It further allowed us to reject low p_t tracks at the trigger level.

The tracking with this apparatus used an array of four cylindrical straw tube drift chambers with charge division, providing us with 3D position information. The chambers were used both for triggering and for final track reconstruction. The signals from two fan shaped hodoscopes of scintillating counters served as input to the first level trigger, which started the digitization of the event [6], [7], [8], [9], [10].

The variable we present in this paper is the *transparency ratio* T_{CH} , which we define to be the ratio of $C(p, 2p)$ quasi-elastic(QE) to pp elastic cross-section divided by the Z of the nucleus(6 in our case) for a restricted region of kinematics near that of pp elastic scattering. Within the impulse approximation this quantity is related to nuclear transparency via

$$T_{CH} = \frac{C}{ZH} = T_r \int_{\alpha_1}^{\alpha_2} \frac{d\alpha}{\alpha} \int \int d\vec{p}_{Ft} n(\alpha, \vec{p}_{Ft}) \frac{\frac{d\sigma}{dt}(s)}{\frac{d\sigma}{dt}(s_0)} . \quad (2)$$

Here $n(\alpha, \vec{p}_{Ft})$ is the momentum distribution of the target proton in the nucleus, $\alpha = \frac{E_F - p_{Fz}}{m} \simeq \frac{s}{s_0}$ near $\alpha = 1$, s and s_0 are the Mandelstam variables for C and H events correspondingly, $\vec{p}_{Ft} = (p_{Fx}, p_{Fy})$ is the transverse component of the target

nucleon momentum, $\frac{d\sigma}{dt}(s)$ is the differential pp cross-section for the process inside the nucleus, $\frac{d\sigma}{dt}(s_0)$ - for the hydrogen.

In the presentation of these measurements, we attempt to limit the influence of incomplete knowledge of the spectral function. This was accomplished by applying an unrestrictive cut on transverse Fermi momentum but a tight cut on longitudinal nuclear motion so that $\frac{d\sigma}{dt}(s) \simeq \frac{d\sigma}{dt}(s_0)$, neglecting small variations in cross-section.

Our apparatus was capable of measuring all three components of momenta of both outgoing particles. Therefore the kinematic variables of interest were calculated from the data. The following cuts ensured that the conditions were satisfied: $|p_{Fy}| < 0.3$ GeV/c, $|p_{Fx}| < 0.5$ GeV/c and $0.95 < \alpha_0 < 1.05$. The variable α_0 , defined as

$$\alpha_0 = 1 - \frac{2p \cos \frac{\theta_1 - \theta_2}{2} \cos \frac{\theta_1 + \theta_2}{2} - p_{inc} z}{m} \quad (3)$$

with $p = \sqrt{(\frac{E_{beam} + m}{2})^2 - m^2}$, is an analog of α , which was defined to be independent of momentum variables and varies only with the opening angle in the Lab frame. The value of $|\alpha - \alpha_0|$ in the region of interest is well below one percent [7], [8].

To extract the number of events from C and CH_2 targets, we used the missing mass variable defined as:

$$M_{miss}^2 = E_{miss}^2 - \vec{p}_F^2 \quad (4)$$

As it follows from equation (4), M_{miss}^2 distribution for the reconstructed QE events should display a peak around zero. From the events that satisfy the cuts we select candidates for QE by demanding exactly two charged tracks in the final state. We also select events with more than two tracks to model the behavior of the background. The signal is then extracted from these distributions after background subtraction.

The hydrogen signal was measured as the normalized difference between the yields from the CH_2 and C targets. Then the transparency ratio can be defined as

$$T_{CH} = \frac{\beta \gamma C}{3(CH_2 - \beta \gamma C)} \quad (5)$$

Here C and CH_2 are the numbers of extracted events. $\beta = 0.4535$ is the ratio of number of CH_2 molecules to the number of C atoms in the corresponding targets, γ is the beam normalization factor $\gamma = \frac{beam_C}{beam_{CH_2}}$. The beam normalization factor is determined from the flux measurements using scaler readings and comparing yields for the same target positions and different target configurations.

We performed our measurements at incoming momenta of 5.9 and 7.5 GeV/c in 1994 [11] and 5.9, 8.0, 9.0, 11.7 and 14.4 GeV/c in 1998. These corresponded to the following values of momentum transfer Q^2 : 4.8 and 6.2 (GeV/c)² in 1994 and 4.8, 6.6, 7.6, 10.1 and 12.7 (GeV/c)² in 1998. The preliminary result of our experiment, the energy dependence of the transparency ratio, integrated over the




FIGURE 1. The measured dependence of the transparency ratio, T_{CH} , on the incident momentum, P_{inc} . Boxes represent 1994 results and circles - 1998. No systematic errors are included.

center-of-mass scattering angle in the region $85^\circ - 90^\circ$, is presented in the figure 1. We can draw the following conclusions from it. First, the ratio definitely varies with energy with strong deviations from Glauber calculations. Second, we clearly see the rise of the ratio for the incident momenta below 10 GeV/c. Third, we also observe the fall of the transparency above 10 GeV/c. Finally, our results seem to be in a good agreement with the energy dependence reported in 1988 by E834 [2]

REFERENCES

1. G.R. Farrar et al., *Phys. Rev. Lett.* **61**, 686 (1988).
2. A.S. Carroll et al., *Phys. Rev. Lett.* **61**, 1698 (1988).
3. B. Pire and J. Ralston, *Phys. Rev. Lett.* **61**, 1823 (1988).
4. C.F. de Teramond and S.J. Brodsky, *Phys. Rev. Lett.* **60**, 1924 (1988).
5. M. Strikman, L. Frankfurt and G.A. Miller, *Comm. Nucl. Part. Phys.* **21**, 1 (1992).
6. J. Wu et al., *Nucl. Instr. and Meth.* **349**, 183 (1994).
7. I. Mardor, *PhD Thesis*, unpublished, Tel-Aviv University, 1997.
8. Y. Mardor, *PhD Thesis*, unpublished, Tel-Aviv University, 1997.
9. S. Durrant, *PhD Thesis*, unpublished, The Pennsylvania State University, 1994.
10. A. Leksanov, *PhD Thesis*, in preparation, The Pennsylvania State University, 2000.
11. I. Mardor et al., *Phys. Rev. Lett.* **81**, 5085 (1998).